

GEOPHELIA TRANQUILLA. *Geop. facie et gutture pallidè cinereis, occipite dorso alisque e cinereo-fuscis, singulis plumis ad apicem angustè nigerrimo fasciatis; alulis spuriiis, primariisque saturatè fuscis, pectore, lateribus, et nuchâ pallidè cinereis, lineis angustis nigris crebrè notatis, abdomine et lateribus pallidè vinaceis, abdomine medio crissoque albis; humeris subtùs castaneis.*

Face and throat pale grey; occiput, back and wings ashy brown, each feather bounded at the end with a narrow band of deep velvety black; spurious wing and primaries dark brown; chest, sides and back of the neck pale grey, crossed by numerous narrow, irregular bands of black; abdomen and flanks pale vinous; centre of the abdomen and under tail-coverts white; under surface of the shoulder deep chestnut; four centre tail-feathers greyish brown, passing into black at the tip; the lateral tail-feathers black, largely tipped with white; irides transparent bluish white; base of bill and nostrils light blue; tip of the bill bluish black; naked skin of the orbits deeply wrinkled and of a beautiful light greenish blue; frontal scales of the tarsi and toes dark purple; hind part of the legs flesh-colour.

Total length, $8\frac{3}{4}$ inches; bill, $\frac{5}{8}$; wing, 4; tail, $4\frac{3}{4}$; tarsi, $\frac{5}{8}$.

Hab. Liverpool plains and banks of the Namoi, interior of New South Wales.