

GLYCIPHILA FASCIATA. *Glyc. vertice fuscescenti-nigro, plumis ad apicem lunulá parvulá albá ornatis; uropygio rufo tincto, faciei lateribus, gulá et corpore subtùs albis, ab angulis oris strigá angustá fuscescenti-nigrá per latera colli ductá; pectore lineis semicircularibus fuscescenti-nigris transversim fasciato-lateribus et crisso luteolis, laterum plumis strigá centrali fuscescenti-nigrá longitudinalitèr notatis.*

Crown of the head brownish black, with a small crescent of white at the extremity of each feather; feathers of the back very dark brown, margined with buffy brown; rump tinged with rufous; wings and tail dark brown, fringed with light brown; sides of the face, throat, and under surface white; from the angle of the mouth down the side of the neck a narrow stripe of brownish black; chest crossed by a number of semicircular brownish black fasciæ; flanks and under tail-coverts buff, the former with a stripe of brownish black down the centre; irides reddish brown; bill greenish grey; feet aurora-red.

Total length,  $4\frac{3}{4}$  inches; bill,  $\frac{5}{8}$ ; wing,  $2\frac{7}{8}$ ; tail,  $2\frac{1}{8}$ ; tarsi,  $\frac{5}{8}$ .

*Hab.* Port Essington.